

Workers who are exposed to extreme cold or work in cold environments may be at risk of cold stress. Extremely cold or wet weather is a dangerous situation that can cause occupational illness and injuries such as hypothermia, frostbite, trench foot, and chilblains.

Hypothermia

A condition in which the body uses up its stored energy and can no longer produce heat. Often occurs after prolonged exposure to cold temperature.

Early symptoms

- Shivering
- Fatigue
- Loss of coordination
- Confusion and disorientation

Late symptoms

- No shivering
- Blue skin
- Dilated pupils
- Slowed pulse and breathing
- Loss of consciousness

First Aid

- Request immediate medical assistance.
- Move the victim into a warm room or shelter.
- Remove wet clothing.
- Warm the center of their body first—chest, neck, head, and groin—using an electric blanket; or use skin-to-skin contact under loose, dry layers of blankets, clothing, or towels.
- If conscious, warm beverages may help increase the body temperature. Do not give alcohol.
- Once temperature has increased keep them dry and wrapped in a warm blanket, including the head and neck.
- If no pulse, begin CPR.

Trench Foot

An injury of the feet resulting from prolonged exposure to wet and cold conditions that can occur at temperatures as high as 60 °F if the feet are constantly wet.

Symptoms

- Reddening of the skin
- Numbness
- Leg cramps
- Swelling
- Tingling pain
- Blisters or ulcers
- Bleeding under the skin
- Gangrene (foot may turn dark purple, blue, or gray)

First Aid

- Remove shoes/boots and wet socks.
- Dry feet.
- Avoid walking on feet, as this may cause tissue damage.

